

SUMMARY OF REQUIREMENTS



MINIMUM AGE AND MINIMUM INTERVALS

This table should be used to determine minimum intervals for “catching up” children who have fallen behind. Otherwise, the recommended schedule should be used.

VACCINE	Minimum AGE Dose 1	Minimum INTERVAL Dose 1 to 2	Minimum INTERVAL Dose 2 to 3	Minimum INTERVAL Dose 3 to 4	Minimum INTERVAL Dose 4 to 5
DTaP/DTP/DT	6 weeks	4 weeks	4 weeks	6 months <i>NOTE: For auditing purposes only — dose 4 need not be repeated if given at least 4 months after dose 3.</i>	6 months
Polio	6 weeks	4 weeks	4 weeks	6 months (final dose must be given on or after the fourth birthday AND 6 months from the previous dose)	
MMR	12 months	4 weeks	If the first dose of MMR is given <u>before</u> the first birthday, it must be repeated.		
Hepatitis B	birth	4 weeks	8 weeks (child must be at least 24 weeks of age)	There must be 16 weeks between dose 1 and 3.	
Hib	6 weeks	4 weeks If first dose given at <12 months 8 weeks (as final dose) If first dose given at 12-14 months No further doses needed If first dose given at ≥15 months	4 weeks If current age <12 months 8 weeks (as final dose) If current age ≥12 months and second dose given at <15 months No further doses needed If previous dose given at ≥15 months	8 weeks (as final dose) This dose only necessary for children aged 12 months - 5 years who received 3 doses before 12 months. Last dose should not be given earlier than 12 months and a minimum of 8 weeks after previous dose.	<i>NOTE: Schedule may vary according to child's current age and previous number of doses received.</i>
Varicella (Chickenpox)	12 months	3 months (Children >13 years of age need 2 doses, 4 weeks apart.)	If the first dose of Varicella is given <u>before</u> the first birthday, it must be repeated.		
Hepatitis A	12 months	6 months			
Pneumococcal Conjugate (PCV)	6 weeks	4 weeks If first dose given at <12 months 8 weeks (as final dose) If first dose given at ≥12 months or current age 24-59 months No further doses needed for healthy children if first dose given at ≥24 months	4 weeks If current age <12 months 8 weeks (as final dose) If current age ≥12 months No further doses needed for healthy children if previous dose given at ≥24 months	8 weeks (as final dose) This dose necessary for children age 12 months - 5 years who received 3 doses before age 12 months.	<i>NOTE: One supplemental dose of Prevnar 13 vaccine is recommended for healthy children 14-59 months of age who have completed the 4-dose series with Prevnar 7 AND for children with underlying medical conditions through 71 months of age. Schedule may vary according to child's current age and previous number of doses received.</i>
Tetanus/Diphtheria/Pertussis (Tdap)	A single dose of Tdap is recommended at 11-12 years of age, regardless of interval since the last tetanus/diphtheria-containing vaccine. Td boosters should be administered every ten years thereafter.				
Meningococcal	11-12 years	8 weeks			